

Papaya -

Basic Set of SIP – 5 & SNP 8 Applications



Soil Improvement Pack

Day 01, 07, 34, 52, 61



Soil Nourishment Pack

Day 91, 116, 141, 166, 191, 216, 241, 270

Day 01, 180- Land Preparation

2 Tones of on farm prepared Vermi-Compost/
Compost



Mulching – Day 30, 120, 180, 250

On farm available green & dry bio-mass &
border crop dressings of Gliricidia, subabhul,
karanj, agasti



Day 30, 60, 90 Soil Application – drip/drench


1 Itr Poshak-G
Supplements main & micro nutrients



Day 07, 52, 88, 7 to 10 Days After Fruit Set Foliar Application

1 Itr Bahar


Early flower initiation & healthy fruit setting,
optimum fruit size & weight



Day 210, 225, 240, 255 & 7 days before harvesting - 3 Soil & 2 Foliar Applications

1 Itr Poshak-K

Supplements main & micro nutrients
Enhances fruit size, weight & color



7 days before harvesting Foliar Application

1 Itr Calplus

Enhances shelf life



Pests & Diseases Control

Major Pest -

Aphid, Mealybug, Stem Borer, White Fly, Fruit Fly

Major Diseases -

Anthracnose, Blights, Leaf Spots, Powdery Mildew, Fruit Rot



Control Remedies -

01. Regular field Check-up for any pest pressure & diseases symptoms. Use colored sticky traps for the same. The application of soil nourishment pack makes the crop pest & disease resistant. SNP application develops useful biology in and around crop which takes care of harmful pests & diseases. SNP also boosts crop immunity. Major pest & disease problems are taken care by SIP & SNP application
02. If observed attack of pest or disease is to considerable extent, then only apply no-synthetic, non-poisonous organic/biological inputs to very limited extent
03. To control pest pressure, use any one of the following input - Neem Oil, Verticillium Lecanii, Beauveria Bassina, Metarhizium Anisoplia, Nomarks
04. To control disease pressure, use any one of the following input – Bordo, Buttermilk, Tricoderma Viride, Pseudomonas Fluorescens, Fungistop
05. Do not apply any input as preventive measure
06. Do not make any soil application of pest or disease controlling input
07. **Crop Rotation** – No mono-cropping pattern to be followed. Crop rotation with proper management of cropping & fallow land is to be practiced. For perennial crops, inter-cropping, mulching to be practiced

01. Farming is a full time business. To be managed accordingly.
02. Border cropping for green manure & wind breaking. Gliricidia, subabul, karanj, agasti, tulas.
03. Field layout for crop plan & rotation scheduleing.
04. Regular green/dry mulching
05. Desi cow/s management
06. Gradual reduction of synthetic chemicals to conclude with no chemical usage
07. Use of Planting Calendar

Disclaimer - The application intimations are based on best of our knowledge. We have tried to collect data from various sources to make the suggestions closer to accuracy. These are indicative intimations; actual requirement may defer based on climatic & crop condition. In any case what so ever, we are not responsible for any compensation & or claims for crop injury, ineffectiveness or other unintended consequences.