

## Tomato -

### Basic Set of SIP - 5 & SNP 4 Applications



Soil Improvement Pack

Day 01, 07, 25, 43, 61



Soil Nourishment Pack

Day 91, 106, 121, 136

### Land Preparation

2 Tones of on farm prepared Vermi-Compost/  
Compost



### Mulching - Day 30 & If Required

On farm available green & dry bio-mass &  
border crop dressings of Gliricidia, subabul,  
karanj, agasti



### Nursery - Day 07, Transplantation - Day 30, 60 Soil Application - drip/drench

500 ml Poshak-G  
Supplements main & micro nutrients



**Nursery – Day 07, Transplantation - 30 & 45  
Foliar + Soil Application**

**500 ml Bahar**

**Early optimum flowering & healthy fruit size, weight**



**Day 50, 61, 72, 83, 94, 105, 116, 127  
250 ml Soil + 250 ml Foliar Application**

**500 ml Poshak-K + 250 ml Calplus**

**Supplements main & micro nutrients Enhances fruit size, weight & color and shelf life**



## Pests & Diseases Control

### Major Pest –

Aphids, Fruit Worm, Leaf Miner, Thrips, Hone Worm

### Major Diseases –

Anthracnose, Canker, Early Blights, Grey Mould, Leaf Curl, Leaf Spot, Leaf Mould



### Control Remedies -

01. Regular field Check-up for any pest pressure & diseases symptoms. Use colored sticky traps for the same. The application of soil nourishment pack makes the crop pest & disease resistant. SNP application develops useful biology in and around crop which takes care of harmful pests & diseases. SNP also boosts crop immunity. Major pest & disease problems are taken care by SIP & SNP application
02. If observed attack of pest or disease is to considerable extent, then only apply non-synthetic, non-poisonous organic/biological inputs to very limited extent
03. To control pest pressure, use any one of the following input - Neem Oil, Verticillium Lecanii, Beauveria Bassina, Metarhizium Anisoplia, Nomarks
04. To control disease pressure, use any one of the following input – Bordo, Buttermilk, Tricoderma Viride, Pseudomonas Fluorescens, Fungistop
05. Do not apply any input as preventive measure
06. Do not make any soil application of pest or disease controlling input
07. **Crop Rotation** – No mono-cropping pattern to be followed. Crop rotation with proper management of cropping & fallow land is to be practiced. For perennial crops, inter-cropping, mulching to be practiced

01. Farming is a full time business. To be managed accordingly.
02. Border cropping for green manure & wind breaking. Gliricidia, subabul, karanj, agasti, tulas.
03. Field layout for crop plan & rotation scheduling.
04. Regular green/dry mulching
05. Desi cow/s management
06. Gradual reduction of synthetic chemicals to conclude with no chemical usage
07. Use of Planting Calendar

Disclaimer - The application intimations are based on best of our knowledge. We have tried to collect data from various sources to make the suggestions closer to accuracy. These are indicative intimations; actual requirement may defer based on climatic & crop condition. In any case what so ever, we are not responsible for any compensation & or claims for crop injury, ineffectiveness or other unintended consequences.